



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The reason for this I attribute to the attempt of the flowers to secure pollination. The plant was kept in a room, and while the flowers were few in number there was no chance for the transfer of pollen, as was easily the case when they became more numerous and crowded. The stamens appeared to wilt in about two days after their pollen had been thrust out by the styles and had they, as was observed in later cases, drawn the styles down into the tube with them, then the object of their living would have been defeated. In two cases I transferred pollen to the stigmas and no movement of the styles was noticed independent of the stamens, but after a time both styles and stamens were drawn down within the tube. My conclusion is that undoubtedly the first cases failed of fertilization and the withdrawing of the styles and the subsequent unfolding of the style branches was a plan to longer present their stigmas for the reception of pollen.—WALTER H. EVANS, *Herbarium Eli Lilly & Co., Indianapolis.*

An abnormal water-pore.—The accompanying figure illustrates a curious water-pore found by Mr. E. L. Hicks, a student in the botanical laboratory of the University of Wisconsin, while examining these



structures on the leaves of *Tropaeolum majus*. The four guard cells bound a somewhat trapezoidal pore, A. The whole apparatus reminds one strikingly of a stoma of *Marchantia polymorpha*. That it was a functionally active pore was shown by the distinct incrustation of the guard-cells with mineral salts.—C. R. B.

A new grass: Melica? multinervosa.—Culms from a strong creeping rhizoma, about 3 ft. high, somewhat

thickened at the base, erect, smooth, frequently geniculate below, the lower nodes hairy: leaves four or five, narrow, rather rigid, 6 to 12 in. long, becoming involute; lower sheaths much longer than the internodes and open above, upper sheaths shorter; ligule a prominent ring of hairs: panicle erect, 6 inches long, the branches single, the lower ones 3 inches long, flowering above the middle with 3 to 6 single, alternate, short-pedicelled, approximate spikelets, the upper branches gradually shorter, above nearly sessile, the lower branches spreading somewhat in flowering; rachis angular, scabrous, hairy in the main axils: spikelets spindle-shaped or linear-lanceolate, 6 to 9 lines long, 8 to 12-flowered, slightly compressed, the flowers imbricated, purple on the margins; empty glumes somewhat unequal, the lower 2 lines long, 1- or faintly 3-nerved, the upper 7-nerved, both